

EXISTENCE OF THREE SOLUTIONS FOR A FIRST-ORDER PROBLEM WITH NONLINEAR NONLOCAL BOUNDARY CONDITIONS

DOUGLAS R. ANDERSON

ABSTRACT. Conditions for the existence of at least three positive solutions to the nonlinear first-order problem with a nonlinear nonlocal boundary condition given by

$$\begin{aligned} y'(t) - p(t)y(t) &= \sum_{i=1}^m f_i(t, y(t)), \quad t \in [0, 1], \\ \lambda y(0) &= y(1) + \sum_{j=1}^n \Phi_j(\tau_j, y(\tau_j)), \quad \tau_j \in [0, 1], \end{aligned}$$

are discussed, for sufficiently large $\lambda > 1$. The Leggett-Williams fixed point theorem is utilized.

1. INTRODUCTION

We are interested in the first-order boundary value problem with nonlinear nonlocal boundary condition given by

$$y'(t) - p(t)y(t) = \sum_{i=1}^m f_i(t, y(t)), \quad t \in [0, 1], \quad (1.1)$$

$$\lambda y(0) = y(1) + \sum_{j=1}^n \Phi_j(\tau_j, y(\tau_j)), \quad \tau_j \in [0, 1], \quad (1.2)$$

where $p : [0, 1] \rightarrow [0, \infty)$ is continuous; the nonlocal points satisfy $0 \leq \tau_1 < \tau_2 < \cdots < \tau_n \leq 1$; the nonlinear functions $\Phi_j : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ satisfy

$$0 \leq y\phi_j(t, y) \leq \Phi_j(t, y) \leq y\psi_j(t, y), \quad t \in [0, 1], \quad y \in [0, \infty), \quad (1.3)$$

for some positive continuous (possibly nonlinear) functions $\phi_j, \psi_j : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$; the scalar λ satisfies

$$\lambda > \left(1 + \sum_{j=1}^n \beta_j\right) \exp\left(\int_0^1 p(\eta) d\eta\right) > 1, \quad \beta_j := \max_{[0, 1] \times [0, C]} \psi_j(t, y) \quad (1.4)$$

for some real constant $C > 0$; and the nonlinear functions $f_i : [0, 1] \times [0, \infty) \rightarrow [0, \infty)$ are all continuous. We also set

$$\alpha_j := \min_{[0, 1] \times [B, \lambda B]} \phi_j(t, y) \quad (1.5)$$

2010 *Mathematics Subject Classification.* 34B05.

Key words and phrases. positive solutions, cone, fixed point theorem, nonlinear boundary condition, Leggett-Williams theorem.

for some constant real $B > 0$ for later reference. Note that by continuity and compactness α_j and β_j exist and satisfy $\beta_j > \alpha_j > 0$.

Some of the motivation for this paper and the study of problem (1.1), (1.2) is as follows. First-order equations with various boundary conditions, including multi-point and nonlocal conditions, are of recent interest. For example, we cite the following papers. Zhao and Sun [14] were concerned with the first-order PBVP (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) + p(t)y(t) = \lambda f(t, y(t)), \quad t \in [0, 1], \quad (1.6)$$

$$y(0) = y(1). \quad (1.7)$$

Tian and Ge [12] investigated the first-order three-point problem (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) + p(t)y(t) = \lambda f(t, y(t)), \quad t \in [0, 1], \quad (1.8)$$

$$y(0) - \alpha y(\eta) = \gamma y(1), \quad (1.9)$$

while Gao and Luo [2] were interested in the problem (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) + p(t)y(t) = \lambda f(t, y(t)), \quad t \in [0, 1], \quad (1.10)$$

$$y(0) = \sum_{j=1}^n \gamma_j y(t_j); \quad (1.11)$$

similarly Anderson [1] studied the first-order problem (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) + p(t)y(t) = \lambda f(t, y(t)), \quad t \in [0, 1], \quad (1.12)$$

$$y(0) = y(1) + \sum_{j=1}^n \gamma_j y(t_j). \quad (1.13)$$

In a related paper, Nieto and R. Rodríguez-López [8] considered

$$y'(t) + p(t)y(t) = \lambda f(t), \quad t \in [0, 1], \quad (1.14)$$

$$\lambda y(t_0) = \sum_{j=1}^n \gamma_j y(t_j). \quad (1.15)$$

Gilbert [3] looked at (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) = \lambda f(t, y(t)), \quad a.e. \ t \in [0, 1], \quad (1.16)$$

$$y(0) = y(1), \quad \text{or} \quad y(0) = y_0 \quad (1.17)$$

using measure theory and Δ -Carathéodory functions. Goodrich [4] analyzed the p -Laplacian problem (if $\mathbb{T} = \mathbb{R}$)

$$\phi_p(y'(t)) = h(t)f(y(t)), \quad t \in [0, 1], \quad (1.18)$$

$$y(0) = \psi(y) \quad \text{or} \quad y(0) = B_0(y'(1)) \quad \text{or} \quad y(0) = (y'(1))^m, \quad (1.19)$$

while Graef and Kong [6] explored the related p -Laplacian problem (if $\mathbb{T} = \mathbb{R}$)

$$\phi_p(y'(t)) = f(t, y(t)), \quad t \in [0, 1], \quad (1.20)$$

$$y(0) = B(y'(1)). \quad (1.21)$$

Otero-Espinar and Vivero [9] gave a general view of different kinds of weak first-order discontinuous boundary value problems on an arbitrary time scale with nonlinear functional boundary value conditions. Shu and Deng [11] proved the existence of three positive solutions to (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) = \lambda f(y(t)), \quad t \in [0, 1], \quad (1.22)$$

$$y(0) = \gamma y(1). \quad (1.23)$$

Zhao [13] applied a monotone iteration method to the problem (if $\mathbb{T} = \mathbb{R}$)

$$y'(t) + p(t)y(t) = \lambda f(t, y(t)), \quad t \in [0, 1], \quad (1.24)$$

$$y(0) = g(y(1)), \quad (1.25)$$

where g denotes a nonlinear boundary condition. Precup and Trif [10] dealt with the existence, localization and multiplicity of positive solutions to non-local problems for first order differential systems of the form

$$y'(t) = f(t, y(t)), \quad t \in [0, 1], \quad (1.26)$$

$$y(0) = \alpha[y], \quad (1.27)$$

where α is linear and continuous. In a second-order problem, Goodrich [5] was concerned with (if $\mathbb{T} = \mathbb{R}$)

$$y^{\Delta\Delta} = -\lambda f(t, y(t)), \quad t \in [0, 1], \quad (1.28)$$

$$y(0) = \phi(y), \quad y(1) = 0. \quad (1.29)$$

where ϕ is a nonlocal boundary condition. One can see from these representative works that problem (1.1) with the nonlinear nonlocal boundary condition (1.2) is new.

2. AT LEAST THREE POSITIVE SOLUTIONS

In this section we establish the existence of at least three positive solutions for the boundary value problem (1.1), (1.2), by finding fixed points for operators on cones in a Banach space; the theorem we will use below is the Leggett-Williams fixed point theorem [7].

A nonempty closed convex set P contained in a real Banach space S is called a cone if it satisfies the following two conditions:

- (i) if $y \in P$ and $\zeta \geq 0$ then $\zeta y \in P$;
- (ii) if $y \in P$ and $-y \in P$ then $y = 0$.

The cone P induces an ordering $\leq$ on S by $x \leq y$ if and only if $y - x \in P$. An operator K is said to be completely continuous if it is continuous and compact (maps bounded sets into relatively compact sets). A map θ is a nonnegative continuous concave functional on P if it satisfies the following conditions:

- (i) $\theta : P \rightarrow [0, \infty)$ is continuous;

(ii) $\theta(tx + (1-t)y) \geq t\theta(x) + (1-t)\theta(y)$ for all $x, y \in P$ and $0 \leq t \leq 1$.

Let

$$P_C := \{y \in P : \|y\| < C\} \quad (2.1)$$

and

$$P(\theta, A, B) := \{y \in P : A \leq \theta(y), \|y\| \leq B\}. \quad (2.2)$$

Theorem 2.1 (Leggett-Williams). *Let P be a cone in the real Banach space S , $K : \overline{P_C} \rightarrow \overline{P_C}$ be completely continuous, and θ be a nonnegative continuous concave functional on P with $\theta(y) \leq \|y\|$ for all $y \in \overline{P_C}$. Suppose there exist constants $0 < A < B < B^\dagger \leq C$ such that the following conditions hold:*

- (i) $\{y \in P(\theta, B, B^\dagger) : \theta(y) > B\} \neq \emptyset$ and $\theta(Ky) > B$ for all $y \in P(\theta, B, B^\dagger)$;
- (ii) $\|Ky\| < A$ for $\|y\| \leq A$;
- (iii) $\theta(Ky) > B$ for $y \in P(\theta, B, C)$ with $\|Ky\| > B^\dagger$.

Then K has at least three fixed points y_1, y_2 , and y_3 in $\overline{P_C}$ satisfying:

$$\|y_1\| < A, \quad \theta(y_2) > B, \quad A < \|y_3\| \text{ with } \theta(y_3) < B.$$

In the next theorem we use the following notation. Let

$$G(t, s) = \frac{\exp\left(\int_s^t p(\eta) d\eta\right)}{\lambda - \exp\left(\int_0^1 p(\eta) d\eta\right)} \times \begin{cases} \lambda & : 0 \leq s < t \leq 1, \\ \exp\left(\int_0^1 p(\eta) d\eta\right) & : 0 \leq t \leq s \leq 1, \end{cases} \quad (2.3)$$

be the corresponding Green's function,

$$M := \frac{1}{\int_0^1 G(1, s) ds} \left(1 - \frac{\exp\left(\int_0^1 p(\eta) d\eta\right) \sum_{j=1}^n \beta_j}{\lambda - \exp\left(\int_0^1 p(\eta) d\eta\right)} \right) > 0 \quad (2.4)$$

for β_j from (1.4), and

$$N := \frac{1}{\int_0^1 G(0, s) ds} \left(1 - \frac{\sum_{j=1}^n \alpha_j}{\lambda - \exp\left(\int_0^1 p(\eta) d\eta\right)} \right) > 0 \quad (2.5)$$

for α_j from (1.5). Both M and N are positive by (1.4).

Theorem 2.2. *Suppose that (1.3) and (1.4) hold, and suppose that there exist constants $0 < A < B < \lambda B \leq C$ such that*

- (F₁) $f_i(t, y) < MA/m$ for $t \in [0, 1]$, $y \in [0, A]$,
- (F₂) $f_i(t, y) > NB/m$ for $t \in [0, 1]$, $y \in [B, \lambda B]$,
- (F₃) $f_i(t, y) \leq MC/m$ for $t \in [0, 1]$, $y \in [0, C]$,

for $i = 1, 2, \dots, m$, where M and N are given above by (2.4) and (2.5), respectively. Then the boundary value problem (1.1), (1.2) has at least three positive solutions y_1, y_2, y_3 satisfying

$$\|y_1\| = y_1(1) < A, \quad B < \theta(y_2) = y_2(0), \quad \|y_3\| = y_3(1) > A \text{ with } \theta(y_3) = y_3(0) < B,$$

where θ is given below in (??).

3. CLOSING COMMENTS

Although we deal with problem (1.1), (1.2) on the real unit interval, the boundary value problem and accompanying techniques introduced in this work can be readily extended to related difference equations and dynamic equations on time scales.

The main conditions here are (1.3) on the nonlinear functions, and on λ in (1.4). Note that these assumptions are fairly mild. We do not assume that the Φ_j are completely separable, nor that they are asymptotically linear. As for λ , we merely have it bounded below, leaving an unbounded range of possible values that it may assume.

Future authors may wish to explore (1.1), (1.2) using monotone iteration methods, upper and lower solution techniques, existence of solutions under singularity, positone, or semi-positone conditions, and so on. We leave the details to interested readers.

REFERENCES

- [1] D. R. Anderson, Existence of solutions for first-order multi-point problems with changing-sign nonlinearity, *J. Difference Equations Appl.*, 14:6 (2008) 657–666.
- [2] C.-H. Gao and H. Luo, Positive solutions to nonlinear first-order nonlocal bvps with parameter on time scales, *Boundary Value Problems*, Volume 2011 (2011), Article ID 198598, 15 pages.
- [3] H. Gilbert, Existence theorems for first-order equations on time scales with Δ -Carathéodory functions, *Advances Difference Equations* Volume 2010 (2010), Article ID 650827, 20 pages.
- [4] C. S. Goodrich, Existence of a positive solution to a first-order p -Laplacian bvp on a time scale, *Nonlinear Anal.*, 74 (2011) 1926–1936.
- [5] C. S. Goodrich, Positive solutions to boundary value problems with nonlinear boundary conditions, *Nonlinear Anal.*, 75 (2012) 417–432.
- [6] J. R. Graef and L. Kong, First-order singular boundary value problems with p -Laplacian on time scales, *J. Difference Equations Appl.*, 17 No. 5 (2011) 831–839.
- [7] R. W. Leggett and L. R. Williams, Multiple positive fixed points of nonlinear operators on ordered Banach spaces, *Indiana U. Math. J.*, 28 (1979) 673–688.
- [8] J. J. Nieto and R. Rodríguez-López, Green's function for first-order multipoint boundary value problems and applications to the existence of solutions with constant sign, *J. Math. Anal. Appl.*, 388 (2012) 952–963.
- [9] V. Otero-Espinar and D.R. Vivero, The existence and approximation of extremal solutions to several first-order discontinuous dynamic equations with nonlinear boundary value conditions, *Nonlinear Anal.*, 68 (2008) 2027–2037.
- [10] R. Precup and D. Trif, Multiple positive solutions of non-local initial value problems for first order differential systems *Nonlinear Anal.*, (2012), doi:10.1016/j.na.2012.06.008
- [11] W. Shu and C. Deng, Three positive solutions of nonlinear first-order boundary value problems on time scales, *International J. Pure Appl. Math.*, 63 No. 2 (2010) 129–136.
- [12] Y. Tian and W.-G. Ge, Existence and uniqueness results for nonlinear first-order three-point boundary value problems on time scales, *Nonlinear Anal.*, 69:9 (2008) 2833–2842.
- [13] Y.-H. Zhao, First-order boundary value problem with nonlinear boundary condition on time scales, *Discrete Dyn. Nature Society*, Vol. 2011 (2011), Article ID 845107, 8 pages.

- [14] Y.-H. Zhao and J.-P. Sun, Monotone iterative technique for first-order nonlinear periodic boundary value problems on time scales, *Advances Difference Equations*, Vol. 2010 (2010), Article ID 620459, 10 pages.

DEPARTMENT OF MATHEMATICS, CONCORDIA COLLEGE, MOORHEAD, MN 56562 USA

E-mail address: `andersod@cord.edu`